

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Clinical Anatomy and Physiology

Journal homepage: <https://www.ijcap.org/>

Original Research Article

Morphometric analysis of proximal femur in Indian population and its implications in Total Hip Arthroplasty

Jaikumar B Contractor^{1*}, Bhavika Umraniya², Praveen Singh³¹Dept. of Anatomy, GMERS Medical College, Panchmahal, Godhra, Gujarat, India²Dept. of Anatomy, GMERS Medical College, Gotri, Vadodra, Gujarat, India³Dept. of Anatomy, Pramukhswami Medical College, Bhaikaka University, Karamsad, Gujarat, India

ARTICLE INFO

Article history:

Received 09-01-2024

Accepted 23-02-2024

Available online 06-05-2024

Keywords:

Hip prosthesis

Artificial implants

Femur head

Femur neck

Femur anthropometry

ABSTRACT

Background: Geometric understanding of the proximal end of femur is significant for functional bipedal erect posture and in pre-operative planning of osteotomy as well as the design and development of implants for THA. Moreover, the anthropological parameters of any bone are determined by genetic and environmental factors such as age, race, gender and lifestyle and this leads to racial variations in the morphological parameters of proximal end of femur owing to lifestyle, physique, applied force and their distribution in India population.

Materials and Methods: The current study was designed to study and compare the morphology of proximal end of femur in Indian population and was conducted on 94 dry human cadaveric bones of unknown age and sex to determine the morphology of the proximal end of femur.

Results: We observed the Femur Length as 426.6 ± 15.82 mm, Femur Neck Length as 3.455 ± 0.378 mm and Neck Shaft Angle as $125.27 \pm 2.54^\circ$ amongst other parameters.

Conclusion: The findings also revealed significant variations in the morphological parameters among different populations, emphasizing the importance of considering racial diversity in the design and selection of implants for THA for improving the success and longevity of hip arthroplasty procedures.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Total Hip Arthroplasty (THA) is a common method of treating hip joint failure occurring due to osteoarthritis, rheumatoid arthritis, osteonecrosis, trauma and bone tumours.¹ Here the geometric understanding of the proximal end of femur is not only significant for functional bipedal erect posture but becomes an essential parameter in pre-operative planning of osteotomy as well as the design and development of implants for THA.² An accurate measurement is vital in selection of the implant to minimize the risk of complications like aseptic loosening, improper load distribution and discomfort resulting from mismatch

as well as to ensure long term success through proper alignment of the prosthesis to be implanted.³

Moreover, the anthropological parameters of any bone is determined by genetic and environmental factors such as age, race, gender and lifestyle^{4,5} and this leads to racial variations in the morphological parameters of proximal end of femur owing to lifestyle, physique, applied force and their distribution in India population. However, currently a range of standard sized femur implants designed from anthropometric data of western population are used for THA leading to discrepancy in regards to the measurements and this non-availability of data for Indian population can lead to complications in the long run.^{6,7}

* Corresponding author.

E-mail address: jaicontractor02@gmail.com (J. B. Contractor).

The current study was designed to study and compare the morphology of proximal end of femur in Indian population with others.

2. Materials and Methods

The current study was conducted at the Department of Anatomy, Pramukhswami Medical College, Karamsad and the other medical colleges of Central Gujarat Region. Around 94 dry human cadaveric bones of unknown age and sex were studied to determine the morphology of the proximal end of femur. All adult dry femur bones without any visible osseous pathologies like tumours, deformities, fractures, trauma were included in the study.

All measurements of the proximal end of femur were taken using Digital sliding Vernier caliper with 0.01 mm precision, goniometer and an Osteometric board. All measurements were taken by a single author to avoid any inter-observer error and each measurement was repeated thrice to avoid any intra observer error. Arithmetic average of the three readings was considered the final reading for the study.

The morphological examination of the proximal end of femur was done in following two steps:^{8,9}

The following parameters were observed:

1. **Femur length (FL):** Femur was placed in a position parallel to the surface of osteometric board by rotating the femur shaft internally, then the distance between the highest point of the femur head to the lowest point of the medial condyle was measured as the femur length.
2. **Femoral head diameter (FHD):** It was measured as the average of the diameter of the femoral head in the cranio caudal axis (distance in a straight line from the upper end to the lower end of the femoral head) and sagittal axis (distance in a straight line from the front end and behind end of the femoral head).
3. **Femoral neck length (FNL):** The length of femoral neck was measured as the distance between the inferior region of base of femoral head and the lower end of intertrochanteric line.
4. **Femoral neck width (FNW):** The diameter of the femoral neck in cranio-caudal axis (cc-axis) was measured as the distance in a straight line from the upper end to the lower end of the anatomical neck of the femur and the diameter of femoral neck in sagittal axis (s-axis) was measured as the distance in a straight line from the front end to the rear end of the femur.
5. **Neck-shaft angle (NSA):** It was measured as the angle intersected between the long axis of the shaft of femur and the long axis of the neck of femur. Femoral shaft axis was considered as a vertical line from the tip of greater trochanter and Femoral neck axis was drawn by joining the center of head of femur and the

midpoint of Intertrochanteric line. NSA was measured in the frontal plane by means of goniometry.

6. **Length of intertrochanteric line (LIL):** It was measured as the distance in a straight line joining the highest and lowest point of trochanters.

The data was tabulated in Microsoft Office Excel 2016 software (Microsoft Corp.). The results were tabulated and analyzed using descriptive statistics. For the statistical analysis the Unpaired Student T-test was used to compare the morphological parameters between population of different geographic locations.

3. Results

The results obtained through morphometric analysis have been depicted in Table 1.

4. Discussion

As observed in Tables 2 and 3 we found a significant difference for the measured value when compared to those of different studies at the given degree of freedom (DF), i.e. 'p' value was < 0.05 implying that the morphometric parameters of proximal femur are significantly different across different populations.

Racial differentiation in the morphometric parameters of proximal femur across different populations have also been reported by De Sousa E et al²⁰ in their study evaluating the variables with Auto CAD 2000 in Brazilian population. Baharuddin MY et al²¹ concluded that femur in Malaysian population were generally smaller and different than western femur in many morphological parameters. Umer et al.²² also reported that the morphology of proximal femur in standardized antero-posterior pelvic radiographs in Pakistani population differed significantly than that from western population. These studies highlight the racial variations in parameters of proximal femur.

According to Reddy et al,²³ an implant mismatch, has been strongly correlated with increased risk of intra-operative fracture or limb lengthening as well as micro-motion leading to increased incidences of anterior thigh pain, improper load distribution, aseptic loosening as well as osteolysis. Implants and prosthesis designed for western population are larger in size as well as the angles and orientations are mismatch to other populations. It emphasizes the need to design these implants based on anthropometric and bio-mechanic data for a specific population thereby minimizing complications.

Since subdivisions of Indian population do not have specific implants designed for them, the observations of present study can be used to replicate the normal anatomy as far as possible. Improved knowledge of the morphometric parameters of proximal femur will not only aid surgeons during total hip arthroplasty but the data could also be used as a guideline to design appropriate fit implants for the

Table 1: Morphometric measurements of proximal femur in mm

	FL	FHD		FNL	FNW		NSA	LIL
		cc-axis	s-axis		cc-axis	s-axis		
Mean ± SD	426.6 ± 15.82	39.60 ± 2.28	37.69 ± 3.99	3.455 ± 0.378	31.60 ± 2.32	26.47 ± 3.20	125.27 ± 2.54	6.71 ± 0.54
Median	42.63	39.36	37.67	3.48	32.09	27.28	125.00	6.70
Range	383.7 – 465.7	35.51 – 45.27	31.16 – 46.85	2.13 – 4.47	23.76 – 41.63	19.31 – 31.51	116 – 132	5.8 – 8.8

Table 2: Comparison of morphometric parameters of femur with Indian population

S. No.	Parameters	Present Study	Ravi et al ¹⁰	Khan SM & Saheb SH ¹¹	Gujjar et al ¹²	Siwach RC ¹³	Dhivya S et al ¹⁴
1.	Population	Western Gujarat	South Indian	South Indian	Western Gujarat	North Indian	South Indian
2.	Sample Size	94	592	250	250	150	158
3.	FL	426.62 ± 15.82	447.1 ± 28.94*	446.2 ± 26.39*	438 ± 25.64*	36.9 ± 4.11*	41.66 ± 3.03*
4.	NL	34.55 ± 3.78	36.3 ± 5.4*	36.3 ± 4.2*	34.4 ± 3.8*	37.23 ± 4.65*	—
5.	NSA	125.27 ± 2.54	136.80 ± 4.45*	137	136.30 ± 6.0*	123.5 ± 4.34*	134.15 ± 5.52*
6.	FHD	39.60 ± 2.28	—	—	—	43.95 ± 3.06*	—
	S-axis	37.69 ± 3.99	—	—	—	—	—
7.	FNW	31.60 ± 2.32	—	—	—	31.87 ± 2.91	—
	S-axis	26.47 ± 3.20	—	—	—	24.90 ± 2.94*	—
8.	LIL	6.71 ± 0.54	—	—	—	—	—

*The p-value is significant when the respective study is compared with the present study. (i.e. p < 0.05)

Table 3: Comparison of morphometric parameters of femur with different population

S. No.	Parameters	Present Study	Menezes TM et al ¹⁵	Zulyan T, Murshid KA ¹⁹	Umanuntana A et al ¹⁶	Cho HJ et al ¹⁷	Rubin PJ et al ¹⁸
1	Population	Indian	Brazil	Arabian	Americans & Caucasians	Korean	French
	Type of measurement	Cadaveric bones	Cadaveric bones	Cadaveric bones	Digital Photographs	3D Reconstruction	Radiographic Reconstruction
2	Sample Size	94	29	36 (R) 36 (L)	200	202	32
3	FL	426.62 ± 15.82	416.8 ± 68.6*	428.4 ± 24.9*	132.69 ± 5.91*	130.27 ± 6.25*	443.6 ± 21.8*
4	NL	34.55 ± 3.78	2.55±0.42*	43.4 ± 3.2*	52.09 ± 4.43*	45.50 ± 3.39*	122.9 ± 7.6*
5	NSA	125.27 ± 2.54	4.42 ± 0.44*	44.3 ± 3.3*	—	—	43.4 ± 2.6*
6	FHD	39.60 ± 2.28	4.38 ± 0.47*	30.7 ± 3.6	—	—	—
	CC-axis	37.69 ± 3.99	3.10 ± 0.35	26.3 ± 3.1	—	—	—
7	FNW	31.60 ± 2.32	2.50 ± 0.37	—	—	—	—
	S-axis	26.47 ± 3.20	4.79 ± 0.62*	—	—	—	—
8	LIL	6.71 ± 0.54	—	—	—	—	—

*The p-value is significant when the respective study is compared with the present study. (i.e. p < 0.05)

population.

Since the study was done on dry cadavers it was not possible to ensure that right and left sided femurs belonged to a particular individual. Moreover, we did not observe significant difference in measurements for right and left sided femur, hence the data has been presented accordingly. We also did not categorise the data for gender, but it will be noteworthy to observe any significant difference there as well as observe the horizontal and vertical offset of the femurs for further understanding.

5. Conclusion

The findings revealed significant variations in the morphological parameters among populations, emphasizing the importance of considering racial diversity in the design and selection of implants for THA for improving the success and longevity of hip arthroplasty procedures in India.

6. Source of Funding

None.


7. Conflict of Interest


None.

References

- Amstutz HC. Complications of total hip replacement. *Clin Orthop Relat Res.* 1970;72:123–37.
- Husmann O, Rubin PJ, Leyvraz PF, Roguin BD, Argenson JN. Three-dimensional morphology of the proximal femur. *J Arthroplasty.* 1997;12(4):444–50.
- Engh CA, Bobyn JD, Glassman AH. Porous-coated hip replacement. The factors governing bone ingrowth, stress shielding, and clinical results. *J Bone Joint Surg Br.* 1987;69(1):45–55.
- Nurzinski MK, Briffa NK, Price RI, Khoo BC, Devine A, Beck TJ, et al. Geometric indices of bone strength are associated with physical activity and dietary calcium intake in healthy older women. *J Bone Miner Res.* 2007;22(3):416–24.
- El-Najjar MY, McWilliams KR. Forensic Anthropology: The structure, morphology and variations human bone and dentition. USA: Charles C Thomas Pub Ltd; 1977.
- Jain AK, Maheshwari AV, Singh MP, Nath S, Bhargav SK. Femoral neck anteversion: A comprehensive Indian study. *Indian J Orthop.* 2005;39:137–44.
- Siwach RC, Dahiya S. Anthropometric Study of Proximal Femur Geometry and Its Clinical Application. *Indian J Orthop.* 2003;37(4):247–51.
- Mourao ALM, De-Vasconcellos HA. Proximal femur geometry in Brazilian bones. *Acta Fisiatr.* 2001;8(3):113–9.
- Murlimanju BV, Prabhu LV, Pai MM, Kumar BM, Dhananjaya KVN, Prashanth KU. Osteometric study of the upper end of femur and its clinical applications. *Eur J Orthop Surg Traumatol.* 2012;22(3):227–30.
- Ravi GO, Saheb SH, Joseph-N AR. A Morphometric Study of Femur and Its Clinical Importance. *Int J Integ Med Sci.* 2016;3(7):341–4.
- Khan SM, Saheb SH. Study on neck shaft angle and femoral length of south indian femurs. *Int J Anat Res.* 2014;2(4):633–5.
- Gujar S, Vikani S, Parmar J, Bondre KV. A Correlation between Femoral Neck Shaft Angle to Femoral Neck Length. *Int J Biomed Adv Res.* 2013;4(5):295–8.
- Siwach R. Anthropometric Study of Proximal Femur Geometry and Its Clinical Application. *Ann Natl Acad Med Sci.* 2018;54(4):203–15.
- Dhivya S, Nandhini V. A Study of Certain Femoral Metrics in South Indian Population and its Clinical Importance. *Int J Sci Study.* 2015;3(7):132–5.
- Menezes TM, Rocha T, De-Oliveira B, De-Albuquerque Y, Caiaffo V. Proximal femoral epiphysis: Manual morphometry versus digital morphometry. *Int J Morphol.* 2015;33(333):1114–9.
- Unnanuntana A, Toogood P, Hart D, Cooperman D, Grant RE. Evaluation of proximal femur geometry using digital photographs. *J Orthop Res.* 2010;28(11):1399–404.
- Cho HJ, Kwak DS, Kim IB. Morphometric evaluation of Korean Femur by geometric computation: comparisons of the sex and the population. *Biomed Res.* 2015;doi:10.1155/2015/730538.
- Rubin PJ, Leyvraz PF, Aubaniac JM, Argenson JN, Esteve P, Roguin B. The morphology of the proximal femur. A three dimensional Radiographic analysis. *J Bone Joint Surg[Br].* 1992;74(1):28–32.
- Zulan T, Murshid KA. An analysis of Anatolian human femur anthropometry. *Turk J Med Sci.* 2002;32:231–5.
- De-Sousa E, Fernandes RMP, Mathias MB, Rodrigues MR, Ambram AJ, Babinski MA. Morphometric study of the proximal femur extremity in Brazilians. *Int J Morphol.* 2010;28(3):835–40.
- Baharuddin MY, Zulkifly AH, His M, Aziz AA. Three dimensional morphometry of the femur to design the total hip Arthroplasty for Malay Population. *Adv Sci Lett.* 2013;19(10):2982–7.
- Umer M, Sepah A, Khan A. Morphology of the proximal femur in the Pakistan population. *J Orthop Surg.* 2010;18:279–81.
- Reddy VS, Moorthy GV, Reddy SG. Do we need a special design of femoral component of total hip prosthesis in our patients? *Indian J Orthop.* 1999;33:282–4.

Author biography

Jaikumar B Contractor, Associate Professor  <https://orcid.org/0000-0003-4915-4097>

Bhavika Umraniya, Assistant Professor  <https://orcid.org/0009-0008-1539-4061>

Praveen Singh, Professor  <https://orcid.org/0000-0002-6145-2954>

Cite this article: Contractor JB, Umraniya B, Singh P. Morphometric analysis of proximal femur in Indian population and its implications in Total Hip Arthroplasty. *Indian J Clin Anat Physiol* 2024;11(1):27–31.