

Effect of Height and BMI on Nerve Conduction Velocity

Amruta S Bennal^{1,*}, Manohar Y Pattar², R H Taklikar³

¹Associate Professor in Physiology, ²Neuropsychitrist, ³Professor and HOD Physiology, Navodaya Medical College, Raichur

Corresponding Author:

E-mail: amrutabennal@gmail.com

ABSTRACT

Objective: To know the effect of Height and BMI on median motor nerve conduction velocity.

Background: Nerve conduction study is part of the electro diagnostic procedures that help in establishing the type and degree of abnormalities of nerve. Nerve conduction velocity is affected by many physiological and technical factors. So we have conducted this study, to know the effect of height and BMI on median motor nerve conduction velocity.

Materials and Methods: A Cross – Sectional study was conducted in 50 female medical students with mean age 19 to 22 years. Nerve conduction velocity was calculated by recording evoked electromyogram (EMG) by stimulating median nerve at elbow and at wrist with the help of EMG electrodes and isolated stimulator by using Power lab 8/30 series with dual Bioamplifier. Statistical analysis was done by using Karl Pearson Correlation coefficient.

Results: There is moderately negative correlation with the height on motor nerve conduction velocity with “r” value -0.23487 and also slight negative correlation with BMI on conduction velocity with “r” value -0.1832.

Conclusion: With increasing height and BMI, nerve conduction velocity is decreased so this phenomenon should be considered for comparative studies and diagnosing pathological conditions.

Keywords: BMI; Height; Nerve Conduction velocity

| Access this article online | |
|-------------------------------------------------------------------------------------|-------------------------------------------|
| Quick Response Code: | Website: www.innovativepublication.com |
|  | DOI: 10.5958/2394-2126.2015.00028.6 |

INTRODUCTION

Electrical conduction of motor and sensory nerves of the human body is assessed by Nerve Conduction Velocity, which is a part of electro-diagnostic procedures, that help in establishing the type and nature of the nerve and commonly used to evaluate function of nerve. Nerve conduction velocity is affected by many physiological and technical variables. Physiological variables such as age, height, gender, upper limb versus lower limb, temperature affects conduction velocity. Diameter and myelination of the nerve fibers also affect nerve conduction velocity.¹

Nerve conduction studies (NCS) are performed to diagnose the disorders of the peripheral nervous system.^{2,3} These enable the clinicians to differentiate the two major groups of peripheral diseases: demyelination and axonal degeneration.⁴ These also help in localizing the site of the lesions.^{5,6}

Height and low body mass index (BMI) have been reported as risk factor for ulnar neuropathy at elbow and high BMI as risk factor for carpal tunnel syndrome.⁷ BMI was also found to have negative correlation with sensory nerve action potential amplitude.⁸ In assessment of diabetic peripheral

neuropathy, BMI is very important factor to be taken into consideration.⁹ Thus, influence of BMI on nerve conduction study is crucial which has to be taken into consideration while interpreting nerve conduction studies.

Peroneal and sural NCV correlated inversely with height and with estimated axonal length, whereas median motor and sensory NCV failed to show any significant relationship to height.¹³ Many studies have been done previously to evaluate the influence of the anthropometric factors such as age, height and body mass index on the nerve velocities.^{10,11,12} However, a majority of these studies were based on the western population. So our study includes to know the effect of height and body mass index (BMI) on median motor nerve conduction velocity.

MATERIAL AND METHODS

Cross – Sectional and Descriptive study was conducted in the Department of Physiology, Navodaya Medical College, Raichur after obtaining clearance certificate from ethical committee. The study group included 50 female medical students with age 19 to 22 years in secretory phase of menstrual phase. (to avoid the influence of hormones) Subject with present history of fever, neurological abnormalities, any limb deformities and history of systemic diseases like diabetes mellitus, hypertension, were excluded. Informed consent was taken. Height and weight measured and BMI was calculated.

BMI was calculated as per formula: Weight (Kg)/ Height (meter)² (Quetelet, s Index).

Instruments used for recording .Nerve conduction velocity was calculated by recording evoked electromyogram (EMG) by stimulating median nerve at elbow and at wrist with the help of EMG electrodes and isolated stimulator by using Power lab 8/30 series with dual Bioamplifier (AD Instruments Australia, Model No.ML870).

After explaining the procedure, the subject were made to lie down on the couch. They were asked to remove jewellerys. Setting with instrument was done. The EMG electrodes were placed on the abductor pollicis brevis muscle. Active electrode was placed on muscle bulk & reference electrode was placed on tendon.¹ The course of right median nerve was traced. The EMG was recorded by stimulating median nerve using isolated stimulator at wrist first and elbow at latter. The distance between two points of stimulation (one at wrist and another at elbow) was measured. The latent period was noted from recording. Difference in the latent period for two stimulation (one at wrist and another at elbow) was calculated.

Nerve conduction velocity was calculated by formula and expressed in meter/second

$$\text{Velocity} = \frac{\text{Distance between stimulation sites (mm)}}{\text{Difference between latencies (ms)}}$$

THE STATISTICAL ANALYSIS

The results were expressed in terms of Mean±SD. Statistical analysis was done by using Karl Pearson Correlation Coefficient. The data was analyzed by using SPSS 17.0 version statistical software. Microsoft Word and Excel have been used to generate graphs, tables etc.

RESULTS

Cross – Sectional and Descriptive study was undertaken in 50 female medical students with age group 19 to 22 years in secretory phase of menstrual phase. All the students belonging to the age group 19 to 22 years. The mean height of the individuals is 1.6±0.6 meter and BMI is 20.46±1.34 kg/m². Median Motor Nerve conduction velocity in right hand is 58.68±2.67 meter/second.

There is moderately negative correlation effect with height on motor nerve conduction velocity with “r” value -0.23487, and it is non significant. There is slight negative correlation effect of BMI on conduction velocity with “r” value -0.1832 and it non significant.


Table 1: Representing following parameters

| Parameter | Mean±sd |
|---------------------------------------|------------|
| Age (years) | 19.34±0.62 |
| Height (in meters) | 1.6±0.6 |
| BMI(kg/m ²) | 20.46±1.34 |
| Nerve conduction velocity (meter/sec) | 56.58±2.67 |


Table 2: Representing Nerve Conduction Velocity with height and BMI

| Parameter | ‘r’ value | ‘p’ value | Significance |
|-------------------------|-----------|-----------|-----------------|
| Height (in meters) | -0.23487 | >0.05 | Non significant |
| BMI(kg/m ²) | -0.1832 | >0.05 | Non significant |

r—Karl Pearson correlation Co-efficient



Graph 1: Representing correlation between Nerve Conduction Velocity with height



Graph 2: Representing correlation between Nerve Conduction Velocity with BMI

DISCUSSION

The study was conducted to know the effect of height and BMI on nerve conduction velocity in right hand in 50 women during their secretory phase of menstrual cycle. Median motor nerve conduction velocity was studied. There is moderately negative correlation effect with height on motor nerve conduction velocity which is non significant. A negative correlation between distal fiber diameter and height may best explain decreased conduction velocity. Distal axonal tapering in the nerves explains the effect. Even Campbell proposed that a decrease in diameter occurs abruptly at a given distance from the cell body.¹⁵ In mature rabbit nerves, Williams found that peripheral motor axon diameter was about half that of ventral spinal nerve root fibers and, despite an

increase in myelin sheath thickness, there was an overall decrease in total fiber diameter.¹⁶ Height-related slowing of nerve conduction velocity was observed in this study.¹⁷ Clinical recognition of this height effect is important, otherwise an individual with mildly slowed peripheral nerve conduction velocity solely related to large stature may be labeled as abnormal.

Even with BMI, same effect has been observed. There is slight negative correlation effect of BMI on conduction velocity which is non significant which may be due to thicker subcutaneous tissue in the person with higher BMI. As the adipose tissue in epineurium may be related to some extent to amount of body Fat,¹⁴ it is reasonable that the amount of such fat may affect the nerve conduction. Our

observation are in agreement with Awang MS et al who observed slowing of conduction velocity (CV) with increasing BMI in median motor nerve.¹⁸

Though the height and BMI affects the median motor nerve conduction velocity, these are not statistically significant. These two factors should be considered while diagnosing pathological conditions, otherwise normal individuals may be diagnosed as abnormal and they will be on unnecessary medication.

CONCLUSION

With increasing height and BMI, nerve conduction velocity is decreased and it is statistically non significant. So this phenomenon should be considered for comparative studies and diagnosing pathological conditions. These two biological factors must be taken into consideration while interpreting nerve conduction studies.

ACKNOWLEDGEMENT

Thanks to all the medical students who were voluntarily involved in study.

REFERENCES:

1. P. K Mishra. J Kalita. Clinical Neurophysiology, 2nd edition. page no 2
2. Preston DC, Shapiro BE. Basic nerve conduction studies. In "Electromyography and Neuromuscular Disorders". Butterworth-Heinemann, 1998, 778.
3. Misulis KE, Head TC. Nerve conduction study and electromyography. In "Essentials of Clinical Neurophysiology" 3rd Ed. Pioli SF (eds). Butterworth-Heinemann 2003;987.
4. Kouyoumdjian JA, zanetta DMT, Monta MPA. Evaluation of age, body mass index and wrist index as risk factors for carpal tunnel syndrome severity. *Muscle Nerve*; 25(1): 93-7.
5. Evans BA and Daube JR. A comparison of three electrodiagnostic methods in diagnosing carpal tunnel syndrome. *Muscle Nerve* 1984; 7: 565
6. Stevens JC: AAEM minimonograph 26: the electrodiagnosis of carpal tunnel syndrome. *Muscle Nerve* 1997; 20: 1477-486.
7. Landau ME, Barher KC, Campbell WW. Effect of body mass index on ulnar nerve conduction velocity, ulnar neuropathy at elbow and carpal tunnel syndrome. *Muscle Nerve* 2005; 32(3): 360-363
8. Hasanzadeh P, Oveisgharan S, Sedighi N, Nafissi S. Effect of skin thickness on sensory nerve action potential amplitude. *Clin Neurophysiol* 2008; 119(8): 1824-1828.
9. Boyraz O, Saracoglu M. The effect of obesity on the assessment of diabetic peripheral neuropathy: a comparison of Michigan patient version test and Michigan physical assessment. *Diabetes Res Clin Pract* 2010; 90(3): 256-260.
10. Hennessey WJ, Falco FJ, Goldberg G, et al. Gender and arm length: influence on nerve conduction parameters in the upper limb. *Arch Phys Med Rehabil* 1994; 75:265-9.
11. S. Saeed, M. Akram. Impact of anthropometric measures on sural nerve conduction in healthy subjects. *J Ayub Med Coll Abbottabad* 2008; 20(4).
12. Rivner MH, Swift TR, Crout BO, Rhodes KP. Toward more rational nerve conduction interpretations: the effect of height. *Muscle Nerve*. 1990 Mar; 13(3):232-9.
13. Soudmand R, Ward LC, Swift TR. Effect of height on nerve conduction velocity. *Neurology*. 1982 Apr; 32(4):407-10.
14. Dumitru D. Nerve conduction studies. In: Dumitru D(ed) : *Electrodiagnostic Medicine* . Philadelphia, Hanley and Belfus. 1995; pp. 111-209
15. Campbell WW, Ward LC, Swift TR: Nerve conduction velocity varies inversely with height. *Muscle Nerve* 1981 ; 4~520-523
16. Williams PL, Wendell-Smith CP: Some additional parametric variations between peripheral nerve fibre populations. *J Anat* 1976;109:505-526
17. Rivner MH, Swift TR, Malik K. Influence of age and height on nerve conduction. *Muscle Nerve*. 2001 Sep;24(9):1134-41.
18. Awang MS, Abdul lah JM, Abdul lah MR, Tharakan J, Prasad A, Husin ZA, Hussin AM, Tahir A, Razak SA. Nerve conduction study among healthy Malays. The influence of age, height, and body mass index on Median, ulnar, common peroneal and sural nerves. *Malaysian Journal of Medical Sciences* 2006; 13(2): 19-23.