



Original Research Article

Assessment of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium, phosphorus in tuberculosis patients

Shaik Mahmood^{1,*}¹Dept. of Physiology, Deccan College of Medical Sciences, Hyderabad, Telengana, India

ARTICLE INFO

Article history:

Received 20-12-2020

Accepted 29-12-2020

Available online 11-01-2021

Keywords:

Serum procalcitonin

Adenosine deaminase

Lactate dehydrogenase

Calcium

Phosphorus and thyroid stimulation hormone

Tuberculosis patients

ABSTRACT

Tuberculosis is an infectious disease which is caused by bacteria. The disease is mostly transmitted from person to person, usually by inhaling bacteria – carrying air droplets. Tuberculosis commonly affects the lungs, but it can also affect any other organ of the body such as brain, intestine, kidneys or the spine. Tuberculosis is one of the most ancient disease of mankind for more than four thousand years. It is a chronic disease caused by Mycobacterium tuberculosis and spreads from person to person through air. The new modalities are very much helpful for the diagnosis and treatment of tuberculosis. Unfortunately people are still suffering with this disease. Worldwide it is among the top ten killer infectious disease secondly to HIV and COVID 19. According to World Health Organisation(WHO) tuberculosis is a global pandemic. The present study shows a very strong significant for the assess of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium, phosphorus and thyroid stimulation hormone in tuberculosis patients.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Tuberculosis is an infectious disease caused by mycobacterium bacteria.^{1,2} Tuberculosis typically affects the lungs, but can also effect the other organs of the body such as brain, intestine, kidneys or spine. Tuberculosis is one of the most ancient disease of mankind for more than four thousand years. It is a chronic disease caused by Mycobacterium tuberculosis and spreads from person to person through air.³⁻⁵ The new modalities are very much helpful for the diagnosis and treatment of tuberculosis, unfortunately people are still suffering with this disease. Worldwide it is among the top ten killer infectious disease secondly to HIV and COVID 19.⁶⁻⁸ According to World Health Organisation(WHO) approximately 1.8 million people die with tuberculosis every year.^{9,10} Tuberculosis is a communicable and treatable disease. The new modalities are very much helpful for the diagnosis and treatment of tuberculosis, unfortunately people are still suffering

with this disease. Worldwide it is among the top ten killer infectious disease secondly to HIV and COVID 19. According to World Health Organisation(WHO) tuberculosis is a global pandemic.¹¹⁻¹³ Procalcitonin (PCT), a 116 amino acid is the prohormone precursor of calcitonin, is expressed primarily in C-cells of the thyroid gland and to a smaller extent in neuroendocrine tissue of other organs, such as lungs and intestines. PCT is a marker of inflammatory response to infection. Adenosine Deaminase Activity (ADA) is a commonly used marker for the diagnosis of tuberculosis pleural effusion.¹⁴ ADA is an enzyme involved in purine metabolism and is needed for the breakdown of adenosine and for the turnover of nucleic acid in tissue. LDH is an intracellular enzyme which catalyses the oxidation of L-lactate to pyruvate, the final step in the metabolic chain of anaerobic glycolysis.¹⁵⁻¹⁷ Serum calcium significantly decreased to hypocalcemic levels and serum phosphorus significantly decreased but was within normophosphatemic limits in pulmonary tuberculosis.¹⁸⁻²⁰ Chemotherapy for tuberculosis managed to raise serum levels of both the ions, with hypocalcemia

* Corresponding author.

E-mail address: mahmood_shaikh2001@yahoo.co.in (S. Mahmood).

still persisting in majority of patients during treatment but getting resolved in a significant percentage of patients at the end of 6 months of treatment. Results indicate the need for calcium and phosphorus supplements in tuberculosis patients during chemotherapy.^{21,22} The present study shows a very strong significant for the assessment of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium, phosphorus and thyroid stimulation hormone in tuberculosis patients.^{23–26}

2. Materials and Methods

A total number of one hundred and fifty positive cases of mycobacterium tuberculosis were taken from the out patients department of pulmonology Owaisi Hospital & Research Centre (a teaching hospital of Deccan College of Medical Sciences, Hyderabad, Telangana State, India) These positive cases were compared with two hundred healthy controls. There was found significant assessment of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium, phosphorus in mycobacterium tuberculosis patients and healthy controls. The biochemical parameters were done in Cobas C311 and MinniVidas at Biochemistry Laboratory, Department of Biochemistry, Owaisi Hospital & Research Centre, Hyderabad, Telangna State India (a teaching hospital & research centre to Deccan College of Medical Sciences, Hyderabad, Telangna State India)

3. Results

The levels of Serum Procalcitonin are significant and raised in Tuberculosis Patients compared to Health Controls as

Serum Procalcitonin(ng/dl) 0.5 ± 0.2 10.0 ± 0.25 $0. < 0.001$.

The levels of Serum adenosine deaminase are significant and raised in Tuberculosis Patients compared to Health Controls as

Serum adenosine deaminase (IU/L) 30.0 ± 5.0 74.06 ± 18.5 < 0.001 .

The levels of Serum Lactate dehydrogenase are significant and raised in Tuberculosis Patients compared to Health Controls as

Serum Lactatedehydrogenase (IU/L) 300.36 ± 28.06 442.94 ± 45.85 < 0.001 .

The levels of Serum Calcium are significant and decreased in Tuberculosis Patients compared to Health Controls as

Serum Calcium (mg/dl) 9.34 ± 0.46 7.72 ± 1.02 < 0.001 .

The levels of Serum Calcium are significant and decreased in Tuberculosis Patients compared to Health Controls as

The levels of Serum Phosphorus are significant and decreased in Tuberculosis Patients compared to Health Controls as

Serum Phosphorus (mg/dl) 3.45 ± 0.46 2.06 ± 0.8 < 0.001 .

4. Discussion

Mycobacterium tuberculosis has been rated as the leading cause of mortality due to an infectious disease.^{27,28} Despite aggressive research conducted on this disease and its mechanism, the question still remains, "how to control the disease"? The presence of reliable diagnostic markers is an important factor contributing to the successful treatment of any disease.^{29,30} Serum Procalcitonin, Adenosindeaminase, Lactate dehydrogenase, Calcium and Phosphorus have been reported as a useful biomarker for diagnosis and prognosis of Tuberculosis.^{31,32} Mycobacterium tuberculosis evades the innate antimicrobial defenses of macrophages by inhibiting the maturation of its phagosome to a bactericidal phagolysosome.³³ Phagosome formation triggers a preprogrammed pathway of maturation into the phagolysosome, a process controlled by Ca²⁺. In the present study, the decreased serum calcium levels in tuberculosis patients indicates a decreased availability of calcium for phagolysosome maturation, decreased efficiency of host antimicrobial activity and hence increased severity of the disease.³⁴

5. Conclusion

Although currently available research does not validate the diagnostic utility of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium and phosphorus in mycobacteriumtuberculosis patients. procalcitonin in tuberculosis patients, results of the present study indicate that measurement of serum procalcitonin, adenosine deaminase, lactate dehydrogenase along with serum calcium and phosphorus could prove as a useful diagnostic marker for the disease. The findings imply that it is imperative to crack the underlying mechanism of increase in adenosine deaminase, lactate dehydrogenase and procalcitonin during bacterial infections (namely- Why bacterial infections induce the PrePCT gene? Is PrePCT preferentially proteolysed to PCT over the proteolysis of PCT to calcitonin, to result in the increased serum PCT and so on) to understand and improve its diagnostic utility. The present study encourages further research to validate the role of serum adenosine deaminase, lactate dehydrogenase PCT-serum calcium and phosphorus combination in differential diagnosis of latent versus active tuberculosis and mild versus severe tuberculosis. It also calls for research at the molecular level on the relative rate of post translational modifications of PrePCT and PCT. This could help us understand the specific situations in which serum PCT adenosine deaminase, lactate dehydrogenaseare increased. Serum Calcium and Serum Phosphorus significantly decreased in bacterial infections

Table 1: Comparison of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium and phosphorus in mycobacterium tuberculosis cases and healthy controls

Parameters	Healthy Controls	Tuberculosis	P values
Procalcitonin(ng/dl)	< 0.5 +_ 0.2	10.0± 0.25	< 0.001
Adenosine deaminase (IU/L)	<30.0 +_ 5.0	74.06 ±18.5	<0.001
Lactatedehydrogenase (IU/L)	300.36 +_ 28.06	442.94 ±45.85	<0.001
Calcium (mg/dl)	9.34 +_ 0.46	7.72 ±1.02	<0.001
Phosphorus (mg/dl)	3.45 +_ 0.46	2.06 ±0.8	<0.001

Mean +_ SD = <0.0001

6. Source of Funding

None.

7. Conflict of Interest

The authors declare no conflict of interest.

References

- Christ-Crain M, Stolz D, Bingisser R, Müller C, Miedinger D, Huber PR, et al. Procalcitonin Guidance of Antibiotic Therapy in Community-acquired Pneumonia. *Am J Respir Crit Care Med.* 2006;174(1):84–93. doi:10.1164/rccm.200512-1922oc.
- Schuetz P, Christ-Crain M, Thomann R. Effect of procalcitonin-based guidelines vs standard guidelines on antibiotic use in lower respiratory tract infections: the ProHOSP randomized controlled trial. *JAMA.* 2009;302:1059–66.
- Boussekey N, Leroy O, Georges H, Devos P, d'Escrivan T, Guery B. Diagnostic and Prognostic Values of Admission Procalcitonin Levels in Community-Acquired Pneumonia in an Intensive Care Unit. *Infect.* 2005;33(4):257–63. doi:10.1007/s15010-005-4096-2.
- Masiá M, Gutiérrez F, Shum C, Padilla S, Navarro JC, Flores E, et al. Usefulness of Procalcitonin Levels in Community-Acquired Pneumonia According to the Patients Outcome Research Team Pneumonia Severity Index. *Chest.* 2005;128(4):2223–9. doi:10.1378/chest.128.4.2223.
- Krüger S, Ewig S, Marre R. Procalcitonin predicts patients at low risk of death from community-acquired pneumonia across all CRB-65 classes. *Eur Respir J.* 2008;31:349–355.
- Menendez R, Cavalcanti M, Reyes S, Mensa J, Martinez R, Marcos MA, et al. Markers of treatment failure in hospitalised community acquired pneumonia. *Thorax.* 2008;63(5):447–52. doi:10.1136/thx.2007.086785.
- Tseng JS, Chan MC, Hsu JY. Procalcitonin is a valuable prognostic marker in ARDS caused by community-acquired pneumonia. *Respirol.* 2008;13:505–9.
- Polzin A, Pletz M, Erbes R, Raffenberg M, Mauch H, Wagner S. Procalcitonin as a diagnostic tool in lower respiratory tract infections and tuberculosis. *Eur Respir J.* 2003;21(6):939–43. doi:10.1183/09031936.03.00055103.
- Interleukin-8 in critically ill patients admitted with suspected sepsis. *Am J Respir Crit Care Med.* 2001;164:396–402.
- Seppä Y, Bloigu A, Honkanen PO, Miettinen L, Syrjälä H. Severity Assessment of Lower Respiratory Tract Infection in Elderly Patients in Primary Care. *Arch Intern Med.* 2001;161(22):2709. doi:10.1001/archinte.161.22.2709.
- The predictive value of serum procalcitonin levels in adult patients with active pulmonary tuberculosis. *Jpn J Infect Dis.* 2006;59:164–7.
- Schluger NW, Rom WN. The Host Immune Response to Tuberculosis. *Am J Respir Crit Care Med.* 1998;157(3):679–91. doi:10.1164/ajrccm.157.3.9708002.
- Moulin F. Procalcitonin in children admitted to hospital with community acquired pneumonia. *Arch Dis Child.* 2001;84(4):332–6. doi:10.1136/adc.84.4.332.
- Nyamande K, Laloo UG. Serum procalcitonin distinguishes CAP due to bacteria, Mycobacterium tuberculosis and PJP. *Int J Tuberc Lung Dis.* 2006;10:510–5.
- Naderi M, Hashemi M, Kouhpayeh H. The status of serum procalcitonin in pulmonary tuberculosis and nontuberculosis pulmonary disease. *J Pak Med Assoc.* 2009;59:647–8.
- Liang QL, Shi HZ, Wang K. Diagnostic accuracy of adenosine deaminase in tuberculous pleurisy: a meta-analysis. *Respir Med.* 2008;102:744–54.
- Lott JA, Nemensanzsky E. Lactate dehydrogenase. In: Lott JA, Wolf PL, editors. *Clinical enzymology, a case oriented approach.* New York: Year Book Medical; 1987. p. 213–44.
- Terpos E, Katodritou E, Roussou M. High serum lactate dehydrogenase adds prognostic value to the international myeloma staging system even in the era of novel agents. *Eur J Haematol.* 2010;85:114–9.
- Fantin VR, St-Pierre J, Leder P. Attenuation of LDH-A expression uncovers a link between glycolysis, mitochondrial physiology, and tumor maintenance. *Cancer Cell.* 2006;9(6):425–34. doi:10.1016/j.ccr.2006.04.023.
- Bierman HR, Hill BR, Reinhardt L. Correlation of serum lactic dehydrogenase activity with the clinical status of patients with cancer, lymphomas, and the leukemias. *Cancer Res.* 1957;17:660–7.
- Porcel JM, Esquerda A, Bielsa S. Diagnostic performance of adenosine deaminase activity in pleural fluid: A single-center experience with over 2100 consecutive patients. *Eur J Intern Med.* 2010;21(5):419–23. doi:10.1016/j.ejim.2010.03.011.
- Boonyagars L, Kiertiburanakul S. Use of adenosine deaminase for the diagnosis of tuberculosis: a review. *J Infect Dis Antimicrob Agents.* 2010;27:111–8.
- Valdés L, José ES, Alvarez D, Valle JM. Adenosine deaminase (ADA) isoenzyme analysis in pleural effusions: diagnostic role, and relevance to the origin of increased ADA in tuberculous pleurisy. *Eur Respir J.* 1996;9(4):747–51. doi:10.1183/09031936.96.09040747.
- Perez-Rodríguez E, Castro DJ. The use of adenosine deaminase isoenzymes in the diagnosis of tuberculous pleuritis. *Curr Opin Pulm Med.* 2000;6:259–66.
- Chan TY, Poon P, Pang J, Swaminathan R, Chan CH, Nisar M, et al. A study of calcium and vitamin D metabolism in Chinese patients with pulmonary tuberculosis. *J Trop Med Hyg.* 1994;97(1):26–30.
- Shirai M, Sato A, Suda T, Shichi I, Yasuda K, Iwata M, et al. Chida Calcium metabolism in tuberculosis. *Kekkaku.* 1990;65(6):415–20.
- Hafiez AA, Abdel-Hafez MA, Salem D, Abdou MA, Helaly AA, Aarag AH. Calcium homeostasis in untreated pulmonary tuberculosis. I-Basic study. *Kekkaku.* 1990;65(5):309–16.
- Ali-Gombe A, Onadeko BO. Serum calcium levels in patients with active pulmonary tuberculosis. *Afr J Med Med Sci.* 1997;26(1-2):67–8.
- Ijaz A, Mehmood T, Saeed W, Qureshi AH, Dilawar M, Anwar M, et al. Calcium abnormalities in pulmonary tuberculosis. *Pak J Med Res.* 2004;4:34.
- Sharma SC. Serum calcium in pulmonary tuberculosis. *Postgrad Med J.* 1981;57(673):694–6. doi:10.1136/pgmj.57.673.694.
- Godwin AO, Johnson DJ, Otimenbhor JO. Total serum calcium and inorganic phosphate levels in tuberculosis patients in benin city Nigeria sierra leone. *J Biomed Res.* 2010;2(2):87–90.

32. Endres DB, Rude RK. Mineral and bone metabolism. In: Tietz Fundamentals of Clinical chemistry. Philadelphia: W.B. Saunders; 2001. p. 795.
33. Baig MA, Mustafa KKG, Lutufullah B, Qureshi MA. Low serum calcium associated with tuberculosis. *Prof Med J.* 2006;13(4):583–6.
34. Rohini K, Srikumar PS, S J, Kumar AM, Bhat S. Assessment of serum calcium, phosphorus, c-reactive protein and procalcitonin in tuberculosis patients. *Int J Collab Res Intern Med Pub Health.* 2012;4(12):1868–75.

Author biography

Shaik Mahmood, Assistant Professor

Cite this article: Mahmood S. Assessment of serum procalcitonin, adenosine deaminase, lactate dehydrogenase, calcium, phosphorus in tuberculosis patients. *Indian J Clin Anat Physiol* 2020;7(4):384-387.