

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Clinical Anatomy and Physiology

Journal homepage: <https://www.ijcap.org/>

Original Research Article

Role of yoga in type 2 diabetes mellitus

Mehak Mahajan^{1,*}, Youshita Mahajan²¹Dept. of Physiology, Government Medical College, Jammu, India²Dept. of ENT, Government Medical College, Jammu, India

ARTICLE INFO

Article history:

Received 27-01-2023

Accepted 10-03-2023

Available online 19-04-2023

Keywords:

FBG

PPBG

T2DM

HbA1c

ABSTRACT

Background: Type 2 Diabetes Mellitus (T2DM) has become most common problem worldwide. Yoga provides best means of self-improvement and gaining full potential of one's body, mind & soul. It has been proved that pranayama and certain Asana's help in lowering blood glucose, balancing blood pressure and calms nervous system. The asanas improve the sensitivity of β -cells to glucose, improves insulin secretion, and improves glucose uptake. Hence, Yogic asanas are important for preventing and curing many ailments. The effect of yogic practices in diabetics has to be investigated well.

Aim & Objectives: To assess the effect of yoga on blood glucose levels and anthropometric parameters before and after doing yoga in T2DM patients.

Materials and Methods: A total of 100 T2DM diabetic patients in age group 30-60 years were taken. Effect of Yoga on fasting Blood glucose (FBG), Post prandial blood glucose (PPBG), HbA1c levels and serum insulin were assessed. Anthropometric parameters like BMI and waist hip ratio were also assessed.

Results: There was significant decrease in FBG, PPBG and HbA1c levels with p value < 0.001 after doing yoga. There was also significant improvement in serum insulin levels and decrease in BMI & waist hip ratio.

Conclusion: A significant improvement in the blood glucose levels occurred in subjects who practiced Yoga. There was also lowering of the drug requirement.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The word yoga means union and aims to balance body, mind, and soul. It is proved that pranayama and certain asanas are important for preventing and curing many diseases. Yoga is also used to achieve good health and cure many diseases. Yoga and pranayama is beneficial on different systems of body. It increases life span and brings balance of psychic and somatic aspect of body.¹⁻⁴ The physiological effects of yoga involve no weight gain, less blood glucose levels, low cholesterol levels and low blood pressure along with improvement in immunity and better

psychological effects.^{5,6}

Type 2 diabetes mellitus is a lifestyle disease caused by insulin resistance with less or absent insulin causing increased blood glucose levels. According to International Diabetes Federation diabetes atlas in 2017, there were roughly 425 million people with diabetes, which will increase to 629 million by 2045.⁷ Sedentary lifestyle and unhealthy dietary habits are major risk factors for lifestyle diseases, like diabetes mellitus. Mental stress also increases the risk and severity of diabetes. Less physical activity increases the risk of diabetes by 3 times and risk of coronary artery disease by 2.4 times.⁸

Yoga is a hope for many diabetic patients to be free from disease and medicines. Yoga involves physical,

* Corresponding author.

E-mail address: mahajanmhk1992@gmail.com (M. Mahajan).

physiological, psychological and endocrinal changes which were reported following yoga.⁹⁻¹¹

Studies have suggested that yoga decreases stress, improves metabolic function, regulates autonomic nervous system and alters hypothalamic-pituitary adrenal axis that releases neural mediators of increased blood sugar.¹²

Hence, the present study was done to assess the role of yoga on blood glucose levels and anthropometric parameters in T2DM patients.

2. Aim & Objectives

To assess the effect of yogic exercises on blood glucose levels i.e., FBG, PPBG, HbA1c and serum insulin levels and to assess effect of yogic exercises on anthropometric parameters i.e., height, weight, BMI and waist-hip ratio in T2DM patients.

3. Materials and Methods

The study population included 100 Type 2 male diabetic patients in age group 30-60 years. Male subjects were taken because they did yoga regularly whereas female subjects didn't do yoga regularly. Hence, follow up was better in male subjects.

3.1. Inclusion criteria

Type 2 diabetic patients.

3.2. Exclusion criteria

Patients with complications like nephropathy, retinopathy and neuropathy. People with liver, pulmonary diseases, thyroid disorder, alcoholics, kidney disease and who were non cooperative.

Patients were explained about the aim of the studies and their consent was taken. They were advised to take healthy diet which included balanced diet and to avoid junk food and processed sugary drinks while patients were also told to continue the same drugs. They were told to perform pranayama and different asanas i.e., Surya Namaskar, Dhanurasana (Bow pose), Paschimottanasana (Seated-forward Bend), Viparita Karani (Legs up the wall), Bhujangasana, Shavasana for a duration of 3 months. After 3 months of yoga, various levels of FBG, PPBG, HbA1c & serum insulin were analysed.

Fasting Blood Glucose was measured by taking blood sample empty stomach by GOD-POD (glucose oxidase peroxidase) method.

Post Prandial Blood Glucose was measured by taking blood sample after meal by GOD-POD method.¹³

HbA1c levels were measured by boronate affinity chromatographic method¹⁴ and serum insulin was measured by ELISA. All these methods were standardized.

They performed Yoga, which included pranayama for 30 min and Yogic Asanas for other 30 min every day. Patients performed Yoga in the Yoga centre under the guidance of yoga instructor.

Height, weight, BMI and waist hip ratio was assessed before and after doing yoga. Body weight was measured without shoes and with light clothing using a mechanical weighing scale. Standing height was measured barefooted with light clothing using a stadiometer. From these parameters, BMI was calculated as per formula: -

$$BMI = \frac{Weight (kg)}{Height (m sq.)}$$

The study was done after taking ethical committee approval & registered vide no: C-312.

3.3. Statistical analysis

Data was presented with the help of appropriate tables. Statistical analysis was done by using SPSS Microsoft version 22. P value < 0.01 was considered as significant.

4. Results

The present study was done in 100 T2DM male subjects. They were subjected to Pranayama and different asanas for a duration of 3 months. There was significant decrease seen in FBG, PPBG & HbA1c levels and serum insulin levels as shown in Table 1 with p value < 0.001.

There was significant decrease in weight, waist hip ratio and BMI with p value < 0.01 as shown in Table 2.

The Table 1 shows the Mean \pm SD of FBG is 152.10 \pm 62.00 & 94.84 \pm 31.40, PPBG is 250.20 \pm 81.60 & 150.02 \pm 44.90, HbA1c is 10.56 \pm 3.42 & 9.02 \pm 3.50, and serum insulin level is 13.9 \pm 4.9 & 9.04 \pm 4.3 of 100 male T2DM patients before and after doing yoga respectively. The difference between them is significant at p value < 0.001.

The Table 2 shows the Mean \pm SD of Height is 168 \pm 2.1, the Mean \pm SD of Weight is 62.90 \pm 2.4 & 61.00 \pm 2.2, BMI is 28.91 \pm 0.9 & 27.00 \pm 0.9 and WHR is 0.90 \pm 0.1 & 0.88 \pm 0.1 of 100 male T2DM patients before and after doing yoga respectively. The difference between them is significant at p value < 0.01 except height that remains the same.

5. Discussion

Fasting blood glucose and post prandial blood glucose levels decreased after yoga in the present study which showed positive effect of yoga in the control of Type 2 Diabetes Mellitus. A significant decrease in these levels has been observed in a study where T2DM patients undergoing yoga who were on hypoglycaemic drugs as compared to those who were only on medication.^{15,16} Similarly, a significant decrease in FBG and PPBG levels was also observed in another study.¹⁷ The beneficial effect of yoga is due to increased insulin sensitivity at target tissues which

Table 1: Shows the effect of yoga on various parameters in diabetic patients

Parameters	Before Yoga	After Yoga	p value
Fasting blood glucose(mg/dl)	152.10 ± 62.00	94.84 ± 31.40	<0.001
Post Prandial blood glucose (mg/dl)	250.20 ± 81.60	150.02 ± 44.90	<0.001
HbA1c (%)	10.56 ± 3.42	9.02 ± 3.50	<0.001
Serum insulin (m IU/ml)	13.9 ± 4.9	9.04 ± 4.3	<0.001

Table 2: Shows the effect of yoga on anthropometric parameters in diabetic patients

Parameters	Before Yoga	After Yoga	p value
Height	168 ± 2.1	168 ± 2.1	
Weight (kg)	62.90 ± 2.4	61.00 ± 2.2	<0.01
BMI (kg/msq.)	28.91 ± 0.9	27.00 ± 0.9	<0.01
Waist Hip Ratio	0.90 ± 0.1	0.88 ± 0.1	<0.01

decreases insulin resistance and improves the peripheral utilization of glucose.¹⁸ It is also proved that yoga can regenerate the beta cells of pancreas.¹⁹ Yoga has good effect on body and decreases stress.²⁰ Thus, for control of T2DM, yoga can be considered as feasible and non-invasive adjuvant therapy. Yoga not only decreases the dosage of oral hypoglycaemic drugs/insulin but also delays the progression of disease and prevents the complications.²¹

A significant decrease in the BMI and waist hip ratio was seen in type 2 DM after yoga. Yoga causes redistribution of fat by causing shift from central obesity to peripheral obesity due to improvement in insulin resistance.²² Another study also showed decrease in weight and even distribution of fat in diabetic patients.²⁰ Chaya et al, showed decrease in waist circumference, weight and BMI in T2DM patients.²³ All these findings are in accordance with our study that has showed significant decrease in weight, BMI, waist circumference, waist hip ratio and improvement in insulin resistance after yoga.

6. Conclusion

The present study showed a significant decrease in fasting and post prandial blood glucose levels in T2DM male patients who practiced Yoga. There was also reduced drug requirement. Yogic practices involving pranayama and different asanas i.e., Surya Namaskar, Dhanurasana (Bow pose), Paschimottanasana (Seated-forward Bend), Viparita Karani (Legs up the wall), Bhujangasana and Shavasana for 3 months decreased body weight, BMI, waist hip ratio and thus improved insulin sensitivity and resistance.

Further, inclusion of adults with T1DM is essential. The psychological burden in T1DM is significantly greater in adults than T2DM using insulin therapy than those not using insulin. Hence, exploring yoga benefits by diabetic type and treatment is required for future research.

7. Source of Funding

None.

8. Conflict of Interest

None.

Acknowledgments

I want to express love and gratitude to my parents Prof (Dr) RK Gupta and Mrs Anju Gupta and special thanks to my elder sister Dr Youshita Mahajan and brother Ishaan Mahajan who inspired me and cooperated with me to work on this topic at all levels. Their affection, love, firm support and perennial encouragement was of immense importance.

I will always remain deeply indebted to my subjects for bearing with me and enriching my experience with each interaction. Above all, I would like to offer my humble gratitude to the Almighty for all the blessing he has showered upon me and will continue to do so throughout my life.


References

1. Madanmohan, Bhavanani AB, Dayanidy G, Sanjay Z, Basavaraddi IV. Effect of Yoga therapy on reaction time, biochemical parameters and wellness score of peri and post-menopausal diabetic patients. *Int J Yoga*. 2012;5(1):10-5.
2. Ankad RB, Herur A, Patil S, Shashikala GV, Chinagudi S. Effect of short-term Pranayama and meditation on cardiovascular functions in healthy individuals. *Heart Views*. 2011;12(2):58-62.
3. Mody BS. Acute effects of Surya Namaskar on the cardiovascular & metabolic system. *J Bodyw Mov Ther*. 2011;15(3):343-7.
4. Pal A, Srivastava N, Tiwari S, Verma NS, Narain VS, Agrawal GG, et al. Effect of Yogic practices on lipid profile and body fat composition in patients of coronary artery disease. *Complement Ther Med*. 2011;19(3):122-7.
5. Telles S, Singh N, Joshi M, Balkrishna A. Posttraumatic stress symptoms and heart rate variability in Bihar flood survivors following Yoga: a randomized controlled study. *BMC Psychiatry*. 2010;10:18.
6. Gopal A, Mondal S, Gandhi A, Arora S, Bhattacharjee J. Effect of integrated Yoga practices on immune responses in examination stress - a preliminary study. *Int J Yoga*. 2011;4(1):26-32.
7. The global picture. In: IDF Diabetes Atlas. vol. 3. International Diabetes Federation; 2017. p. 40-59.
8. Thangasami SR, Chandani AL, Thangasami S. Emphasis of yoga in the management of diabetes. *J Diabetes Metab*. 2015;6(10):613.
9. Udupa KN, Singh RH. The scientific basis of Yoga. *J Amer Med Assoc*. 1972;220:1365.

10. Patel C, North WR. Randomised control trial of Yoga and biofeedback in management of hypertension. *Lancet*. 1975;19(7925):93–5.
11. Jain SC, Uppal A, Bhatnagar SO, Talukdar B. A study of response pattern of noninsulin dependent diabetes to yoga therapy. *Diabetes Res Clin Pract*. 1993;19(1):69–74.
12. Mahajan AS. Role of yoga in hormonal homeostasis. *Int J Clin Exp Physiol*. 2014;1(3):173–8.
13. Sharma SP, Anjankar AP, Kale A. Comparison of glucose levels using glucometer and GOD-POD Method in diabetic patients. *Int J Clin Biochem Resch*. 2017;4(1):6–10.
14. Mallia AK, Hermanson GT, Krohn RI, Fujimoto EK, Smith PK. Preparation and use of a boronic acid affinity support for separation and quantitation of glycosylated haemoglobin. *Anal Lett*. 1981;14:649–61.
15. Amita S, Prabhakar S, Manoj I, Harminder S, Pavan T. Effect of yoga-nidra on blood glucose level in diabetic patients. *Indian J Physiol Pharmacol*. 2009;53(1):97–101.
16. Kumar K. A study on the effect of yogic intervention on serum glucose level on diabetics. *Int J Yoga Allied Sci*. 2012;1(1):68–72.
17. Rajan. Effect of paschimottānāsana and avāsana on serum glucose level of diabetic patients. *Glob J Multidiscip Stud*. 2014;3:191–6.
18. Sahay BK. Role of yoga in diabetes. *J Assoc Physicians India*. 2007;55:121–6.
19. Sahay BK, Murthy KJR. Long-term follow up studies on effect of yoga in diabetes. *Diab Res Clin Pract*. 1988;5(1):S655.
20. Malhotra V, Singh S, Sharma SB, Gupta P, Prasad A, Tandon OP, et al. The status of NIDDM patients after yoga asanas: Assessment of important parameters. *J Clin Diag Res*. 2010;4(3):2652–67.
21. Singh S, Singh KP, Tandon OP, Madhu SV. Influence of pranayamas and yoga-asanas on serum insulin, blood glucose and lipid profile in type 2 diabetes. *Indian J Clin Biochem*. 2008;23(4):365–8.
22. Sahay BK. Yoga and Diabetes. *J Assoc Physicians India*. 1986;34(9):645–8.
23. Chaya MS, Ramakrishnan G, Shastry S, Kishore RP, Nagendra H, Nagarathna R, et al. Insulin sensitivity and cardiac autonomic function in young male practitioners of yoga. *Natl Med J India*. 2008;21(5):217–21.

Author biography

Mehak Mahajan, Registrar

Youshita Mahajan, Research Scholar  <https://orcid.org/0000-0001-8158-4834>

Cite this article: Mahajan M, Mahajan Y. Role of yoga in type 2 diabetes mellitus. *Indian J Clin Anat Physiol* 2023;10(1):6-9.