

The study of gross and CT scan anatomy of lateral nasal wall, infundibulum and sinus drainage pathways and their clinical implications in Government Medical College and Super Facility Hospital, Azamgarh, UP

MK Gupta¹, Manisha Upadhyay^{2*}, Ganesh Kumar³, Anand Bihari⁴

¹Assistant Professor, Dept. of ENT, ²Associate Professor, Dept. of Anatomy, ³Professor, Dept. of Radiology, ⁴Lecturer, Dept. of SPM, Govt. Medical College, Azamgarh, Uttar Pradesh

***Corresponding Author:**

Email: manishaup73@gmail.com

Abstract

Anatomy of lateral nasal wall and infundibulum with special reference to uncinate process is very intricate and variable. The aim of this study was to evaluate the anatomical changes infundibulum, frontal recess and uncinate process and its clinical implications. Changes in anatomy of uncinate process alter the sinus drainage. An endoscopic gross and ct scan study of infundibulum and frontal recess was done on the 50 patients of inflamed sinuses. The results were analysed statistically. Uncinate process attachment to lamina paparacea was found most commonly.

Keywords: Lateral nasal wall, Infundibulum, Inflamed sinuses, Drainage pathways.

Received: 9th August, 2017

Accepted: 27th September, 2017

Introduction


The anatomy of infundibulum and frontal recess alter significantly with changes in anatomy of uncinate process which forms the medial boundary of frontal recess.⁽¹⁾ Uncinate process is a boomerang shaped bone attaching anteriorly to lacrimal bone. It medially covers the maxillary and frontal sinus ostium. Uncinate process bone is covered medially and laterally on both surfaces by mucosa. Removal of uncinate process is the first step in endoscopic sinus surgery. A comprehensive knowledge of anatomy of uncinate process and their anomalies is a must prior to surgery to avoid orbital complications.⁽²⁾ A parsons window is created in lower segment of uncinate process to open the infundibulum. Then with the help of debrider upper and lower portion is removed.^(3,4) Superior attachment of uncinate process is found to differ in different patients.

The anatomical alterations of uncinate process were defined as following by Stammberger and Bolger.⁽⁵⁾

1. Superior attachment of uncinate process are of three types.


A-type I uncinate: Superior attachment to lamina paparacea. Frontal sinus drains in middle meatus.

Type-I




B-type II uncinate: Superior attachment to skull base. Frontal sinus drains in infundibulum.

Type - II



C-Type III uncinate process- Superior attachment to middle turbinate. Frontal sinus drains in ethmoidal infundibulum.



Sometimes the superior end of the uncinate process may be branched.

2. Medially bent uncinate process.
3. Laterally bent uncinate process
4. Hypertrophied uncinate process

In a study conducted on 800 cases, Earwaker^(6,7) provide a detailed description of the variants of superior insertion of uncinate process, by classifying them in association with other variants of osteomeatal complex (ethmoid bulla, middle turbinate, septal deviation, degree of angulation of uncinate process). When the uncinate process is inserted into papyracea lamina, maxillary sinus drainage may be affected.

Materials and Method

This study was carried out in department of otorhinolaryngology and anatomy of GMC Azamgarh, UP from November 2013 to November 2015. A prospective CT scan study was done on 50 patients of chronic rhinosinusitis. CT scan of sinuses were done in coronal and axial view.

Patients with symptoms of nasal congestion, nasaldischarge, headache, facial pain and hyposmia, who were not responding to 3 weeks of medical treatment were evaluated with CT scan paranasal sinuses in coronal view and axial view.

Patients with history of previous sinus surgery were excluded from this study. The data are entry in Cs-Pro software and after that its transfer in SPSS data. Statistical analysis was done using CHI square statistical test with statistical programme for social science version 16.0. A P value <0.05 was considered statistically significant.

Observation

The present study was carried out in department of otorhinolaryngology, Government Medical College, Azamgarh, on 50 patients (100 sides) of chronic sinusitis were included.

Table 1: Sex Distribution

Sex	Percentage %
Male	48
Female	52

Table 2: Age Distribution

Age Group	Percentage %
0-20	18
20-30	24
31-40	44
>40	14

Table 3: Distribution of patient on the basis of different type of bent

Variation	Percentage
Medial Bent	20
Lateral Bent	04
Hypertrophied	08
Pneumatised	08

Table 4: Distribution on the basis of superior attachment

Type of superior attachment	Percentage
Type 1 attached to lamina papyracea	40
Type 2 attached to Skull base	32
Type 3 attached to middle turbinate	18
Type 4 uncinate lying free	10




Fig. 1: Type 1 uncinate process attached to lamina papyracea. Type 3 uncinate Process attached to middle turbinate




Fig. 2: Type 2 uncinate process attached to skull base. Type 4 uncinate process lying free

In this study 52% patients were female and 48% patients were male. Majority of patients, 44% were in age group of 31-40 years. 14% patients belonged to the age above 40 years. 18% belonged to age less than 20 years. 24% patients were in age group of 20 to 30 years. Present study shows that medially bent uncinate was most common (20%). In 4% cases uncinate was found laterally bent. In 8% cases hypertrophied uncinate was found. Pneumatized uncinate was found in 8% cases in this study. Out of 100 uncinate process, type 1 superior attachment was found in 40% patients, type 2 in 32% patients, type 3 in 18% patients and type 4 in 10% patients. Type 1 superior attachment was most common. Few studies had described deviation of uncinate process either medially or laterally leading to narrowing of the infundibulum, frontal and anterior ethmoidal recess producing impaired sinus ventilation in maxillary, frontal and ethmoidal sinuses,^(9,10,11,12) contradicting claims by some studies that deviations of uncinate process prevents contaminated air entering the sinuses.^(13,14,15)

Discussion

Stamberger and Wolf (1988) 16 has documented the preeminent role of osteomeatal complex in causation of chronic rhinosinusitis. Computerized tomographic imaging of osteomeatal complex is of paramount importance in surgery of sinuses. Variations in anatomy of osteomeatal complex are numerous and their detection before surgery is very much significant to avoid complications. Endoscopic examination of nose in conjunction with CT scan paranasal sinuses is the gold standard for the treatment of chronic

rhinosinusitis now-a-days. CT scan of paranasal sinuses accurately detects the bony and soft tissue anatomy of sinuses and their variations. This study was conducted on 50 patients of chronic rhinosinusitis, in which 52% patients were female and 48% male. Majority of patients were in age group of 31 to 40 years (44%).

Uncinate process variations: Uncinate process shows an arc shaped course, therefore in anterior coronal sections the uncinate is wide. In middle third, uncinate lies adjacent to nasolacrimal duct, posteriorly, it is narrow. Free edge of uncinate process may deviate medially. Laterally or anteriorly.⁽¹⁶⁾ Medially bent uncinate process is the most frequent pathological finding in chronic rhinosinusitis patients (Stamberger and Wolf 16). Medially bent uncinate comes in contact with middle turbinate, leading to impaired drainage of paranasal sinuses. In our study, we observed medially bent uncinate process in 20% cases. Laterally bent uncinate process was seen in 4% cases. Hypertrophied uncinate process was seen in 8% cases and pneumatized uncinate in 8% cases. Zinreich⁽¹⁷⁾ noted pneumatization of uncinate process in one patient (0.4%) among 230 patients of chronic rhinosinusitis. Pneumatized uncinate can also be referred to as uncinate bulla. Pneumatization can also impair the sinus drainage. A comparative chart of our findings and findings of other authors is given below. Type three attachment was least common finding. Type one was most common finding.^(18,19,20) Type two finding was in between.

Study	Our study (%)	Tuli et al. (%)	Krzeski et al (%)	Min et al. (%)	Landsberg and Friedman
Type I	40%	79.8	17.83	54	52%
Type II	32%	14	33.12	24.5	-
Type III	18%	3	14.33	21.5	-

A study of Landsberg and Freidman⁽²³⁾ had classified the superior attachment of uncinata process as follows:

Type 1: Insertion of lamina papyracea (LP)

Type 2: Insertion into posterior wall of Agger nassi cell (ANP)

Type 3: Insertion into lamina papyracea and junction of middle turbinate with cribriform plate (MTCP)

Type 4: Insertion into junction of middle turbinate with the cribriform plate

Type 5: Insertion into the ethmoid skull base (ESB)

Type 6: insertion into middle turbinate.

Conclusion

Almost all chronic sinusitis are associated with anatomical variations that alter ventilation. So the preoperative evaluation of variation of uncinata process and its pneumatization helps to avoid intraoperative damage to surrounding structures that alter normal ventilation. From this study we concluded that chronic rhinosinusitis associated with variations in uncinata process. In our study it was found that superior attachment of uncinata process to lamina papyracea Type 1 was most common finding that alters the drainage resulting in frontal sinusitis. Medially bent uncinata process variation was also most common finding. CT scan study is essential before undergoing for functional endoscopic sinus surgery. It provides a precious information regarding variations in uncinata process anatomy, which is helpful in removal of complete disease as well as preventing complications during surgery.

Funding

This study was funded by Government medical college, Azamgarh.

Conflict of interest

Author A and B declared that they have no conflict of interest.

Ethical approval

Article does not contain any study with human or animal participants.

Informed consent

Informed consent was obtained from all the individuals who participated in this study.

References

1. Mafee MF Endoscopic sinus surgery: Role of Radiologist. Am. J. Slnhneuroradiol. 1991;855-60.
2. Earwaker J- Anatomical variation in sinonasal CT. Radio 1993,13:381-415.
3. Zienrich S.J. Albayram S. Benson M. Er al.-The Osteomeatal Complex and Functional Endoscopic Surgery. In Som Pm. Curtin Hd, Editor Head and Neck imaging. 4thEd.St Louis: Mosby.2003;149-74.
4. Bolger WE; Kennedy DW, Bolger WE, Zinreich J(2001) : Anatomy of paranasal sinuses Disease of sinuses, Diagnosis and management. B.C. Decker.
5. Stammberger HR, Bolger WE. Paranasalsinuses, Anatomic terminology and nomenclature. The anatomic terminology group Ann Otol Rhinol Laryngosuppl 1995;167:7-16.
6. Arslan H., Aydinlioglu A., Bozkurt M., et al - Anatomic variations of the paranasal sinuses: CT examination for endoscopic sinus surgery. Auris Nasus Larynx., 1999;26:39-48.
7. Mclaughlin RB. Rehi R.M., Lanza D.C- Clinically Relevant frontal sinus Anatomy And Physiology Otolaryngol Clin North Am.,2001;(34):1-22.380.
8. Bolger WE, Butzin CA, parson DS. - Paranasal sinus bony anatomic variations and mucosal abnormalities: CT analysis for endoscopic sinus surgery. Laryngoscope1991;101.
9. Zinreich SJ, Kennedy DW, Rosenbaum AE, Gayler BW, Kumar AJ, Stamberger H. Paranasal sinuses .CT scan imaging requirement for endoscopic surgery. Radiology 1987;163:769-775.
10. Stammberger H. Secretion transport. In: Functional endoscopic sinus surgery Philadelphia: BC Decker,1991:17-46.
11. Kopp W, Stammberger H, Fötter R. Special radiologic image of the paranasal sinuses. Eur J Radiol 1988;8:152-156.
12. Anita Armani, R.N. Karadi, Saurabhkumar. A study of anatomical variations of Osteomeatal complex in chronic sinusitis patients –CT findings. 2014;8(10):KC)1-KC04.
13. Bolger WE, Woodruff W, Parsons DS. CT demonstration of pneumatization of the uncinata process .Am J Neuroradiol. 1990;11(3):552.
14. Groves J. and Gray R.F. Applied physiology of the nose and paranasal sinuses. In a Synopsis of Otolaryngology.4th edition. John Wright and Sons, Bristol, pp 1995:167-168.
15. Stamberger H, Wolf G Headache and sinus disease. The endoscopic approach Ann Otol Rhinol Laryngosuppl 1988;134,3-23.nomenclature.
16. Tuli IP, Sengupta S, Munjal S, Kesari SP: Anatomical variations of uncinata process observed in chronic sinusitis. Indian J Otolaryngol Head Neck Surg 2013 65(2):15.
17. Mamatha H. Shamsundar NM, Bharathi MB, Prasanna LC (2010) Variations of Osteomeatal complex and Its applied anatomy : a CT Sci Technol 3(8):904–907.
18. Wanamaker HH (1996) Role of Haller's cell in headache and sinus disease: a case report Otolaryngol Head Neck surg 114(2):324-327, doi 10.1016/S0194-5998(96)70196-1.
19. Krzeski A, Tomaszewska E, Jakubczyk I, Galewicz Zielinska A. Anatomic variations of the lateral nasal wall in the computed tomography scans of patients with chronic rhinosinusitis Am J Rhinol 2001;15(6):371-375.
20. Min Y, Koh T, Rhee C, Han M. Clinical implications of the uncinata process in paranasal sinusitis: radiological evaluation. Am J Rhinol.1995;9(3):131-135.
21. Landsberg R, Friedman M. A computer- assisted anatomical study of the nasofrontal region. Laryngoscope. 2001;111:2125-2130.